

ACUTE TOXICITY SUMMARY

CARBON TETRACHLORIDE

(carbon chloride; carbon tet; freon 10; halon-104; methane tetrachloride; necatrine; tetrachlorocarbon; tetrachloromethane; tetraform; tetrasol; univerm)

CAS Registry Number: 56-23-5

I. Acute Toxicity Summary (for a 7-hour exposure)

<i>Inhalation reference exposure level</i>	1,900 µg/m³
<i>Critical effect(s)</i>	toxicity to the developing fetus
<i>Hazard Index target(s)</i>	Reproductive/developmental; Nervous System; Alimentary Tract

II. Physical and Chemical Properties (HSDB, 1994 except as noted)

<i>Description</i>	colorless liquid
<i>Molecular formula</i>	CCl ₄
<i>Molecular weight</i>	153.24
<i>Density</i>	1.59 g/cm ³ @ 20°C
<i>Boiling point</i>	76.54°C
<i>Melting point</i>	-23°C
<i>Vapor pressure</i>	91.3 mm Hg @ 20°C
<i>Flashpoint</i>	critical temperature = 283.1°C
<i>Explosive limits</i>	not found
<i>Solubility</i>	soluble in acetone, ethanol, benzene, carbon disulfide; moderately soluble in water
<i>Odor threshold</i>	96 ppm (604 mg/m ³) (Amoore and Hautala, 1983)
<i>Odor description</i>	sweet, chloroform-like odor
<i>Metabolites</i>	chloroform; carbene radical, carbon monoxide (Ahr <i>et al.</i> , 1980)
<i>Conversion factor</i>	1 ppm = 6.3 mg/m ³

III. Major Uses or Sources

Carbon tetrachloride is used as a solvent for the recovery of tin in tin-plating waste and in the manufacture of semiconductors. It is also used in petrol additives, refrigerants, metal degreasing, and as a catalyst in the production of polymers. Carbon tetrachloride is also used as a chemical intermediate in the production of fluorocarbons and pesticides (HSDB, 1994).

IV. Acute Toxicity to Humans

Hepatotoxicity is the most sensitive and best studied toxic endpoint for CCl₄ exposure (Andrews and Snyder, 1991). The human data on hepatic effects of CCl₄ are based on numerous clinical case reports with poorly defined exposure conditions. The hepatotoxic effects, which may occur more readily in persons regularly consuming alcohol, are often reversible over the course of several weeks (Fry *et al.*, 1959).

Bioactivation of CCl₄ into reactive metabolites by hepatic cytochrome P-450 enzymes results in hepatic centrilobular degeneration and necrosis (Andrews and Snyder, 1991). After a single dose of CCl₄, evidence of centrilobular necrosis is visible within 12 hours and obvious necrosis occurs by 24 hours. If no further injury occurs, the lesions begin to repair after 24 hours, and may be restored to normal after 14 days recovery. A reduction of P450 activity in the liver also occurs and is due to irreversible binding by reactive metabolites and subsequent inhibition of the P450 enzymes that metabolize CCl₄ (Andrews and Snyder, 1991).

Mucosal irritation and CNS effects have also been reported following CCl₄ exposure. In one of the first controlled human studies on the effects of CCl₄, Davis (1934) observed headaches, nausea, and vomiting in subjects exposed to 317 ppm (1,997 mg/m³) for 30 minutes. Stewart *et al.* (1961) exposed 6 human volunteers to 49 ppm (309 mg/m³) CCl₄ for 70 minutes, or to 10-11 ppm (63-69 mg/m³) CCl₄ for 3 hours. The subjects reported no irritation to the eyes or respiratory tract. A Romberg test and a heel to toe test (tests of central nervous system function) were normal in these subjects immediately following exposure, but one individual had elevated urine urobilinogen levels 7 days following exposure. The magnitude of the elevation of urinary urobilinogen was not given. Two of 4 individuals exposed to 49 ppm (309 mg/m³) CCl₄ also exhibited decreased serum iron, although in one the decrease was still within the normal range.

Predisposing Conditions for CCl₄ Toxicity

Medical: Individuals with compromised liver function may be more susceptible to CCl₄-induced hepatotoxicity.

Chemical: Co-exposure to ethanol, acetone, or isopropanol is known to potentiate the toxicity of carbon tetrachloride (Charbonneau *et al.*, 1986; Cornish and Adefuin, 1966). Exposure to other chlorinated compounds, such as chlordecone, also potentiates the toxicity of CCl₄ (Curtis *et al.*, 1979).

V. Acute Toxicity to Laboratory Animals

A concentration of 7,300 ppm (45,990 mg/m³) CCl₄ was reported to be lethal to 1 out of 10 rats after a single, 2-hour exposure (Adams *et al.*, 1952). In this study, one rat out of 30 died after a 10-hour exposure to 3,000 ppm (18,900 mg/m³). Delayed effects from these exposures included weight loss, abnormal behavior and appearance, and additional mortality. Rats surviving these exposures exhibited liver injury evidenced by serum phosphatase, increased prothrombin clotting time, fatty degeneration, and enlargement of the liver.

A 4-hour inhalation exposure to 250 ppm (1,575 mg/m³) CCl₄ resulted in increased serum glutamic-oxalacetic transaminase (SGOT) activity in rats, indicative of hepatic damage (Cornish and Block, 1960). Exposure of these rats to a concentration of 100 ppm for 4 hours did not result in changes in SGOT activity.

Kim and coworkers (1990) showed that administration of a range of concentrations from 10 mg/kg to 1,000 mg/kg CCl₄ by gavage to rats resulted in a dose-dependent increase in serum levels of hepatic enzymes, decrease in hepatic cytochrome P-450 activity, and an increase in centrilobular lesions in the liver. In this study, 10 mg/kg was the LOAEL for hepatocellular changes and elevated serum enzymes. At low doses, the hepatocellular effects of CCl₄ are reversible, while at higher doses necrosis and irreversible damage occur (Gerhard *et al.*, 1970).

In addition to its hepatocellular toxicity, CCl₄ also has been shown to affect the immune system. Mice exposed orally to 500 mg/kg CCl₄ exhibited suppressed T-cell dependent immune responses as measured by decreased splenic antibody forming cells. These mice also had elevated plasma interleukin-2 and transforming growth factor- β 1 measured 24 and 48 hours after exposure (Delaney *et al.*, 1994). However, a previous study showed that hepatotoxicity from CCl₄ occurs at much lower concentrations than does toxicity to the immune system (Smialowitz *et al.*, 1991).

A physiologically-based pharmacokinetic model for carbon tetrachloride has been developed in the rat (Paustenbach *et al.*, 1988). In this model, it was estimated that 60% of inhaled CCl₄ is metabolized, and that 96% of the metabolized CCl₄ forms biological adducts which degrade slowly with a half-life of 24 hours. The remaining 4% of the metabolized CCl₄ becomes CO₂.

VI. Reproductive or Developmental Toxicity

No studies were available on the reproductive effects of CCl₄ in humans. Significant decreases in fetal body weight, crown-rump length, and ossification of sternebrae were observed in rats exposed to 300 ppm (1,890 mg/m³) CCl₄ on days 6-15 gestation (Schwetz *et al.*, 1974). High doses (0.3 ml/100 g body weight) of CCl₄ injected intraventricularly caused marked histologic injury to the chorionic epithelium of the placenta in rats (Tsirel'nikov and Tsirel'nikova, 1976). Carbon tetrachloride has not been listed as a developmental toxicant under Proposition 65.

VII. Derivation of Acute Reference Exposure Level and Other Severity Levels (for a 1-hour exposure)

Mild Adverse Effect Level

Because the most sensitive endpoint found in the inhalation toxicity literature was developmental toxicity, a potentially disabling effect, there is no mild adverse effect level available for CCl₄.

**Reference Exposure Level (protective against severe adverse effects for a 7 hour exposure):
0.3 ppm (1,900 µg/m³)**

Because of the uncertainty in extrapolating from repeated dose studies to a one-hour concentration, for the reproductive/developmental endpoint, we have chosen to use a single day exposure as the basis for the REL. Thus, the REL for CCl₄ is for a 7 hour exposure.

<i>Study</i>	Schwetz <i>et al.</i> (1974)
<i>Study population</i>	pregnant rats
<i>Exposure method</i>	inhalation exposure to 0, 300, or 1,000 ppm for 7 hours/day on days 6-15 of gestation
<i>Critical effects</i>	fetal growth retardation (decreased crown-rump length and body weight)
<i>LOAEL</i>	300 ppm
<i>NOAEL</i>	not determined in this study
<i>Exposure duration</i>	7 hours/day
<i>LOAEL uncertainty factor</i>	10
<i>Interspecies uncertainty factor</i>	10
<i>Intraspecies uncertainty factor</i>	10
<i>Cumulative uncertainty factor</i>	1,000
<i>Reference Exposure Level</i>	0.3 ppm (1.9 mg/m³; 1,900 µg/m³)

Level Protective Against Life-threatening Effects

Rats (5-30 per group, males and females) were exposed to single concentrations of CCl₄ for durations varying from 6 minutes to 10 hours (Adams *et al.*, 1952). Concentrations used ranged from 3,000 to 19,000 ppm (45,990 to 119,700 mg/m³) CCl₄. Mortality was measured for several weeks following the exposure.

1-Hour Mortality Data in Rats from CCl₄ Inhalation

Concentration (ppm x 10 ³)	7.3	8.4	9.4	10.8	12.0	12.0	15.4	17.8	19.0	24.0
Response	0/20	0/20	1/10	1/10	3/10	4/10	7/10	8/10	9/19	20/20

Adams et al. (1952)

A benchmark dose approach used a log-normal probit analysis (Crump, 1983) of rat lethality data from Adams *et al.* (1952). Exposure durations from 1-4 hours were included in the analysis. Concentrations of CCl₄ were adjusted to approximate equivalent 1-hour concentrations using the equation $C^n \cdot T = K$, where $n = 2.8$ (ten Berge *et al.*, 1986). The concentration associated with a 5% incidence of lethality was 8,557 ppm (53,909 mg/m³); the benchmark concentration (BC₀₅) for this response, the 95% lower confidence limit on this concentration, was 7,010 ppm

Determination of Acute Reference Exposure Levels for Airborne Toxicants
March 1999

(44,163 mg/m³). An uncertainty factor (UF) of 3 was applied to the BC₀₅ to account for interspecies variation since the BC₀₅ accounts for some degree of individual variation and a UF of 10 was used to account for human individual variability. The total UF was 30.

$$\text{level protective against life-threatening effects} = \text{BC}_{05} / (\text{UF})$$

The level protective against life-threatening effects for CCl₄ is therefore 234 ppm (1,470 mg/m³). The maximum likelihood estimates (MLE) and 95% lower confidence limits (LCL) for 1% and 5% response rates are compared below. Refer to section IX of this toxicity summary for the graphic representation of benchmark concentration derivation.

Response rate	MLE (ppm)	95% LCL (ppm)
1%	6,646	4,976
5%	8,557	7,010

NIOSH (1995) lists a revised IDLH of 200 ppm based on acute inhalation toxicity in humans.

VIII. References

Adams EM, Spencer HC, Rowe VK, McCollister DD, Irish DD. Vapor toxicity of carbon tetrachloride determined by experiments on laboratory animals. Arch Ind Hyg Occup Med 1952;6:50-66.

Ahr HJ, King LJ, Nastainczyk W, Ullrich V. The mechanism of chloroform and carbon monoxide formation from carbon tetrachloride by microsomal cytochrome P-450. Biochem Pharmacol 1980;29:2855-2861.

Amoore JE, Hautala E. Odor as an aid to chemical safety: Odor thresholds compared with threshold limit values and volatilities for 214 industrial chemicals in air and water dilution. J Appl Toxicol 1983;3(6):272-290.

Andrews LS, Snyder R. Toxic effects of solvents and vapors. In: Amdur MO, Doull J, Klaassen CD, editors. Cassarett and Doull's Toxicology. 4th ed. New York: McGraw Hill; 1991. p. 693-694.

Charbonneau M, Brodeur J, Souich P, Plaa GL. Correlation between acetone-potentiated CCl₄-induced liver injury and blood concentrations after inhalation or oral administration. Toxicol Appl Pharmacol 1986;84:286-294.

Cornish HH, Adefuin J. Ethanol potentiation of halogenated aliphatic solvent toxicity. Am Ind Hyg Assoc 1966;27:57-61.

Cornish HH, Block WD. A study of carbon tetrachloride. I The effect of carbon tetrachloride inhalation on rat serum enzymes. Arch Ind Health 1960;21:549-554.

Determination of Acute Reference Exposure Levels for Airborne Toxicants
March 1999

Crump KS and Co, Inc. Probit (Log-Normal) software for the IBM-PC. Ruston (LA): KS Crump & Co.; 1983.

Curtis LR, Williams WL, Mehendale HM. Potentiation of the hepatotoxicity of carbon tetrachloride following preexposure to chlordecone (Kepone) in the male rat. *Toxicol Appl Pharmacol* 1979;51:283-293.

Davis P. Carbon tetrachloride as an industrial hazard. *JAMA* 1934;103:962-966.

Delaney B, Strom SC, Collins S, Kaminski NE. Carbon tetrachloride suppresses T-cell-dependent immune responses by induction of transforming growth factor- β 1. *Toxicol Appl Pharmacol* 1994;126:98-107.

Fry WA, Smith JM, Suker JR. Acute carbon tetrachloride intoxication. *Northwestern University Med School Quarterly Bulletin* 1959;33:346-351.

Gerhard H, Schultze B, Maurer W. Autoradiographische Untersuchung uber die Proteinsynthesestörung in der Mauseleber nach CCl₄-Intoxikation. *Virchows Arch Abt B Zellpath* 1970;6:38-56.

(HSDB) Hazardous Substances Data Bank. National Library of Medicine, Bethesda (MD) (CD-ROM version). Denver (CO): Micromedex, Inc.; 1994. (Edition expires 1/31/94).

Kim HJ, Odend'hal S, Bruckner JV. Effect of oral dosing vehicles on the acute hepatotoxicity of carbon tetrachloride in rats. *Toxicol Appl Pharmacol* 1990;102:34-49.

(NIOSH) National Institute for Occupational Safety and Health. Chemical listing and documentation of revised IDLH values. 1995. (<http://www.cdc.gov/niosh/intridl4.html>)

Paustenbach DJ, Clewell HJ, Gargas ML, Andersen ME. A physiologically-based pharmacokinetic model for inhaled carbon tetrachloride. *Toxicol Appl Pharmacol* 1988;96:191-211.

Schwetz BA, Leong BK, Gehring PJ. Embryo- and fetotoxicity of inhaled carbon tetrachloride, 1,1-dichloroethane and methyl ethyl ketone in rats. *Toxicol Appl Pharmacol* 1974;28:452-464.

Smialowitz RJ, Simmons JE, Luebke RW, Allis JW. Immunotoxicologic assessment of subacute exposure of rats to carbon tetrachloride with comparison to hepatotoxicity and nephrotoxicity. *Fundam Appl Toxicol* 1991;17:186-196.

Stewart RD, Gay HH, Erley DS, Hake CL, Peterson JE. Human exposure to carbon tetrachloride vapor. *J Occup Med* 1961;3:586-590.

Ten Berge WF, Zwart A, Appelman LM. Concentration-time mortality response relationship of irritant and systemically acting vapours and gases. *J Hazard Mater* 1986;13:301-309.

Determination of Acute Reference Exposure Levels for Airborne Toxicants
March 1999

Tsirel'nikov NI, Tsirel'nikova TG. Morphohistochemical study of the rat placenta after exposure to carbon tetrachloride at different stages of pregnancy. Bull Exp Biol Med 1976;82(1):1262-1267.

U.S. Environmental Protection Agency. Interim methods for development of inhalation Reference Concentrations. EPA 600/8-90/066A. Environmental Criteria and Assessment Office, Office of Research and Development. Cincinnati (OH): U.S.EPA; 1990.